



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07C 217/12, A61P 25/22, A61K 31/13		A2	(11) International Publication Number: WO 00/68183
			(43) International Publication Date: 16 November 2000 (16.11.00)
(21) International Application Number: PCT/HU00/00044		János [HU/HU]; Reviczky Ezredes utca 8, H-1033 Budapest (HU).	
(22) International Filing Date: 10 May 2000 (10.05.00)		(74) Agent: ADVOPATENT OFFICE OF PATENT AND TRADE-MARK ATTORNEYS; P.O. Box 11, H-1251 Budapest (HU).	
(30) Priority Data: P 9901559 11 May 1999 (11.05.99) HU		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(71) Applicant (for all designated States except US): EGIS GYÓGYSZERGYÁR RT. [HU/HU]; Keresztúri út 30-38, H-1106 Budapest (HU).		Published Without international search report and to be republished upon receipt of that report.	
(72) Inventors; and (75) Inventors/Applicants (for US only): LUKÁCS, Gyula [HU/HU]; Bronz u. 5, H-1163 Budapest (HU). SIMIG, Gyula [HU/HU]; Hollósy Simon u. 25, H-1126 Budapest (HU). MEZEI, Tibor [HU/HU]; Borz u. 4, H-1221 Budapest (HU). BUDAI, Zoltán [HU/HU]; Lukács u.3, H-1023 Budapest (HU). PORCS-MAKKAY, Márta [HU/HU]; Lukács Gy. u. 21, H-1039 Budapest (HU). KRASZNAI, György [HU/HU]; XIII u. 38, H-1172 Budapest (HU). NAGY, Kálmán [HU/HU]; Túrista u. 2/a, H-1025 Budapest (HU). VERECZKEYNÉ DONÁTH, Györgyi [HU/HU]; San Marco u. 52, H-1034 Budapest (HU). SZABÓ, Tibor [HU/HU]; Szentmihályi u. 24/c, H-1144 Budapest (HU). NÉMETH, Norbert [HU/HU]; Bartók B. út 92-94, H-1113 Budapest (HU). SZULÁGYI,			
(54) Title: HIGH PURITY (1R,2S,4R)-(-)-2- [(2'-{N,N- DIMETHYLAMINO} -ETHOXY)] -2-[PHENYL] -1,7,7-TRI-[METHYL] -BICYCLO[2.2.1] HEPTANE AND PHARMACEUTICALLY ACCEPTABLE ACID ADDITION SALTS THEREOF AND A PROCESS FOR THE PREPARATION OF THESE COMPOUNDS AS WELL AS MEDICAMENTS CONTAINING 1 OR MORE OF THESE COMPOUNDS AND THEIR USE			
(57) Abstract The invention relates to high purity (1R,2S,4R)-(-)-2- [(2'-{N,N- dimethylamino} -ethoxy)] -2-[phenyl] -1,7,7-tri-[methyl] -bicyclo[2.2.1] heptane and pharmaceutically acceptable acid addition salts thereof containing not more than 0.2 % of (1R,3S,4R)-3- [(2'-{N,N- dimethylamino} -ethyl)]-1,7,7-tri- [methyl] -bicyclo [2.2.1] heptane-2-one and/or of a pharmaceutically acceptable acid addition salt thereof. Furthermore the invention is concerned with a process for the preparation of these compounds. Moreover the invention relates to medicaments containing 1 or more of these compounds and their use.			

FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

High purity (1R,2S,4R)-(-)-2-[(2'-
-{N,N-dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-
-tri-[methyl]-bicyclo[2.2.1]heptane and
pharmaceutically acceptable acid addition
5 salts thereof and a process for the preparation
of these compounds as well as medicaments
containing 1 or more of these compounds and
their use

10 The invention relates to high purity (1R,2S,4R)-(-)-2-
-[(2'-{N,N-dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-tri-
-methyl-bicyclo[2.2.1]heptane and pharmaceutically
acceptable acid addition salts thereof and a process for
the preparation of these compounds as well as medicaments
15 containing 1 or more of these compounds and their use.

The 2-(E)-butenedioate (1 : 1) salt (fumarate) of
(1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-
-phenyl]-1,7,7-tri-methyl-bicyclo[2.2.1]heptane of
20 Formula



I


30 is a known anxiolytic active principle having the INN
"deramciclone fumarate".

The compound of Formula I falls under the general
35 Formula I of Hungarian patent No. 179,164 but has not been
actually and explicitly disclosed in this patent
specification, nor the preparation thereof has been

- 2 -

exemplified. According to Hungarian patent No. 179,164 the alkanol amine cycloalkyl ethers of its general Formula I are prepared by reacting (+)-1,7,7-tri-[methyl]-
-bicyclo[2.2.1]heptane-2-one, i. e. (+)-camphor of Formula

5



II

10


with the corresponding organic metal compound, subjecting the adduct obtained to hydrolysis and introducing onto the hydroxy group of the product obtained the basic side chain by etherification. As organic metal compound a Grignard compound or an organic alkali metal compound, preferably lithium or sodium compound, is used.

20

The preparation of the compound of Formula I has been actually disclosed in Hungarian patent No. 212,574. The essence of this process is that purification of the product is carried out at a later stage of the synthesis. According to the process (+)-camphor of Formula II is subjected to Grignard reaction with phenyl magnesium bromide in diethyl ether to give (1R,2S,4R)-(-)-2-[phenyl]-1,7,7-tri-[methyl]-
-bicyclo[2.2.1]heptane-2-ol of Formula

25

30



III

35

- 3 -

with a yield of 28% (according to GC). The compound (1R,2S,4R)-(-)-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-2-ol of Formula III is in the reaction mixture and is not isolated. The complex is decomposed, the reaction mixture is converted without purification into the sodium salt by reaction with sodium amide or sodium hydride and the sodium salt obtained is reacted with anhydrous (2-{chloro}-ethyl)-dimethylamine in toluene as medium. The reaction mixture contains beside the base (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane of Formula I (being present in an amount of 20 to 30% a considerable amount of impurities and starting materials, e.g. unreacted (+)-camphor of Formula II, (1R,2S,4R)-(-)-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-2-ol, 1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-2-ol and biphenyl, triphenyl impurities, etc. The base (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane of Formula I is separated from said contaminations by extraction with aqueous tartaric acid, whereupon the base is set free and the fumarate salt is formed. The total amount of unreacted (+)-camphor of Formula II and (1R,2S,4R)-(-)-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-2-ol of Formula III remains in the organic phase of the tartaric acid extraction step, which can be re-used in the Grignard reaction after removing the solvent and water (i.e. it can be re-circulated into the process). Thus the (+)-camphor used can be more efficiently utilized; without re-circulation only about 16% by weight of the (+)-camphor used can be utilized, while in case of a one-fold and three-fold re-circulation this value is increased to 22% by weight and 25% by weight, respectively.

It is very important and is to be emphasized that a considerable part of (+)-camphor of Formula II used in the

- 4 -

Grignard reaction does not react and this starting material cannot be technically removed from the desired product because of the physical properties of (+)-camphor and the lability of the compound (1R,2S,4R)-(-)-2-[phenyl]-1,7,7-tri-
5 -[methyl]-bicyclo[2.2.1]heptane-2-ol of Formula III formed since compound (1R,2S,4R)-(-)-2-[phenyl]-1,7,7-tri-
- [methyl]-bicyclo[2.2.1]heptane-2-ol of Formula III is susceptible to decomposition. For this reason according to the process disclosed in Hungarian patent No. 212,574 the
10 alkylation step always takes place in the presence of (+)-
-camphor of Formula II.

The aforesaid gives rise to the drawbacks of the process disclosed in Hungarian patent No. 212,574. The
15 alkali hydrides and amides used in the first step of the alkylation reaction form salts not only with the alcohol (1R,2S,4R)-(-)-2-[phenyl]-1,7,7-tri-[methyl]-
-bicyclo[2.2.1]heptane-2-ol of Formula III but also with (+)-camphor of Formula II and other compounds containing an
20 active hydrogen atom being present in the reaction mixture. For this reason beside the desired compound (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-tri-
- [methyl]-bicyclo[2.2.1]heptane of Formula I further alkylated derivatives formed, e.g. from unreacted
25 (+)-camphor, are obtained and the desired compound (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-
- [phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane of Formula I is to be recovered from a mixture containing such
impurities and also unreacted compounds (+)-camphor and
30 (1R,2S,4R)-(-)-2-[phenyl]-1,7,7-tri-[methyl]-
-bicyclo[2.2.1]heptane-2-ol of Formulae II and III. The crude compound (1R,2S,4R)-(-)-2-[(2'-
- {N,N-dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-tri-
- [methyl]-bicyclo[2.2.1]heptane of Formula I can only be
35 purified, though incompletely, by means of recrystallization from dimethyl formamide. However, with the aid of said recrystallization only non-basic


- 5 -

contaminations can be completely removed, which do not form salts.

A further disadvantage of recrystallization from
5 dimethyl formamide is that the traces of the solvent cannot
be removed from the desired pharmaceutical active principle
to the required extent. In this regard it is to be noted
that according to ICH (International analytical require-
ments accepted by the US, Japan and the EU) the limit of
10 dimethyl formamide is 880 ppm (0.088% by weight). The
reason for that the dimethyl formamide cannot be removed to
such an extent but a greater amount of it remains in the
product is the high boiling point of dimethyl formamide, on
the one hand, and the sensitivity of (1R,2S,4R)-(-)-2-[(2'-
15 -{N,N-dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-tri-
-methyl]-bicyclo[2.2.1]heptane of Formula I to thermal
treatment, on the other hand.

It has been found that in case of the alkylation
20 reaction of (1R,2S,4R)-(-)-2-[phenyl]-1,7,7-tri-methyl-
-bicyclo[2.2.1]heptane-2-ol of Formula III carried out with
(2-{chloro}-ethyl)-dimethylamine (+)-camphor of Formula II
being always present gives rise to the formation of
considerable amounts of by-products, e.g. (1R,3S,4R)-3-
25 -[(2'-{N,N-dimethylamino}-ethyl)]-1,7,7-tri-methyl-
-bicyclo[2.2.1]heptane-2-one of Formula

30



35

- 6 -

The by-product (1R,3S,4R)-3-[(2'-{N,N-dimethylamino}-ethyl)]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-2-one of Formula V is formed as follows: under the conditions used in the etherification reaction (+)-camphor of Formula II forms an alkali salt in position 3 which in turn reacts with the (2-{chloro}-ethyl)-dimethylamine used as alkylating agent to yield the compound of Formula V. The amount of the by-product of Formula V may be as high as 1 to 10%.

The solubility of the fumarate 2-(E)-butenedioate (1 : 1) of the compound of (1R,3S,4R)-3-[(2'-{N,N-dimethylamino}-ethyl)]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-2-one of Formula V is approximately identical with that of the fumarate of the desired compound (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane of Formula I and therefore crystallizes together with the fumarate of the compound (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane of Formula I and contaminates the desired end product. If the etherification is carried out in toluene, as described in Hungarian patent No. 212,574, the product obtained after salt formation in ethanol contains considerable amounts of the impurity (1R,3S,4R)-3-[(2'-{N,N-dimethylamino}-ethyl)]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-2-one of Formula V.

The salt is a highly unsoluble compound and can be recrystallized only from dimethyl formamide. However, recrystallization from dimethyl formamide fails to provide a compound (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane of Formula I in a purity required by the Pharmacopoeias for the following reasons:

- a) The product obtained after recrystallization from dimethyl formamide still contains the compound (1R,3S,4R)-3-[(2'-{N,N-dimethylamino}-ethyl)]-

- 7 -

-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-2-one
of Formula V in an amount above the threshold
value permitted by Pharmacopoeia (about 0.5%);

5 b) Dimethyl formamide has a high boiling point and cannot be removed from the product in the required degree because at the high temperature decomposition of the product takes place.

10 A purification to yield products with a purity
sufficient for medicaments according to the Pharmacopoeias
could not be attained by known purification processes, such
as recrystallization from solvents or fractional
distillation. More specifically, by known processes the
15 (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-
-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane of
Formula I could not be obtained having no more than 0.2
of (1R,3S,4R)-3-[(2'-{N,N-dimethylamino}-ethyl)]-
-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-2-one of Formula
20 V but only with more than 0.5% of this impurity.


Taking into consideration the severe requirements of Pharmacopoeia, impurities being present in an amount higher than 0.2% by weight may endanger the use of the active principle for pharmaceutical purposes. The impurity (1R,3S,4R)-3-[(2'-{N,N-dimethylamino}-ethyl)]-1,7,7-tri-
-[methyl]-bicyclo[2.2.1]heptane-2-one of Formula V may therefore cause problems in the use of the compound of Formula I as active principle.

30 As a summary, it can be stated that when purifying the
(1R,2S,4R) - (-) - 2 - [(2' - {N,N-dimethylamino} - ethoxy)] - 2 -
- [phenyl] - 1,7,7-tri - [methyl] - bicyclo[2.2.1]heptane of
Formula I prepared by the known method, dimethyl formamide
35 would be the only conceivable solvent. However, this
recrystallization method is unsuitable for the preparation
of a pharmaceutical active ingredient meeting the

- 8 -

requirements of the Pharmacopoeias, because dimethyl
formamide has such a high boiling point that traces thereof
cannot be removed from the product to a sufficient extent.
At the high temperature required the compound
5 (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-
- [phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane of
Formula I is subject to decomposition.

As already disclosed above, (+)-camphor of Formula II
10 is present during the alkylation reaction. From (+)-camphor
as further contamination (1R,4R)-2-[(2'-{N,N-
-dimethylamino}-ethoxy)]-1,7,7-tri-[methyl]-
-bicyclo[2.2.1]heptane of Formula



is formed. If an alkali metal hydride or alkali metal amide
is used as basic salt forming agent, the amount of the
contamination (1R,4R)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-
30 -1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane of Formula IV is
1 to 10% The compound (1R,4R)-2-[(2'-{N,N-
-dimethylamino}-ethoxy)]-1,7,7-tri-[methyl]-
-bicyclo[2.2.1]heptane of Formula IV has been known from
the Prior Art [Yakugaku Zasshi, 75, 1377, (1955); Chem.
35 Abstr. 9340 (1956)]. The compound (1R,4R)-2-[(2'-{N,N-
-dimethylamino}-ethoxy)]-1,7,7-tri-[methyl]-
-bicyclo[2.2.1]heptane of Formula IV is formed as follows:

- 9 -

the alkali metal hydride or alkali metal amide used for sodium salt formation in the etherification step reduces 1 to 10% of (+)-camphor of Formula II to borneol which is converted under the reaction conditions used into the alkali metal salt and said alkali salt enters with (2-{chloro}-ethyl)-dimethylamine into an alkylation reaction. However, the borneol ether (1R,4R)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane of Formula IV can be separated from the desired compound (1R,2S,4R)-(-)-2-[(2'-[N,N-dimethylamino]-ethoxy)]-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane of Formula I in the course of working up the reaction mixture.

The problem underlying to the invention is to provide (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane of Formula I and pharmaceutically acceptable acid addition salts thereof containing amounts of (1R,3S,4R)-3-[(2'-{N,N-dimethylamino}-ethyl)]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-2-one of Formula V small enough to meet the requirements of use for pharmaceutical purposes and a process for preparing such compounds without the necessity of recrystallization purification steps which anyhow would lead only to an insufficient purification and would reduce the yield and moreover would have the drawback that the residual solvent could not be removed from the end product to a sufficient extent even by complicated methods, as well as medicaments containing 1 or more of these compounds and their use.

Surprisingly the above has been solved by the present invention which provides a new product that could not be prepared by the Prior Art.


- 10 -

The present invention is based on the surprising recognition that if the reaction between the reaction mixture containing the compound (1R,2S,4R)-(-)-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-2-ol of Formula III and (2-{chloro}-ethyl)-dimethylamine is carried out in the presence of an alkali metal hydride or alkali metal amide in a medium containing dioxane as solvent, the reaction is directed by far in favour of the formation of the desired (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane of Formula I and the by-product (1R,3S,4R)-3-[(2'-{N,N-dimethylamino}-ethyl)]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-2-one of Formula V is formed only in a minimal amount. The above recognition enables the preparation of the desired compound (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane of Formula I containing less than 0.2% of compound (1R,3S,4R)-3-[(2'-{N,N-dimethylamino}-ethyl)]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-2-one of Formula V. The compound (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane of Formula I thus obtained directly meets the requirements of Pharmacopoeia as regards the purity and the content of residual solvent.


In the entire text the percentages regarding the contents of the compounds of Formulae I and V and of other compounds are the result of gas chromatographic analysis they being the ratio of the area under the given peak and the total area under all the peaks.

Hence a subject matter of the invention are (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane of Formula

- 11 -



and pharmaceutically acceptable acid addition salts
15 thereof, characterised by that they contain not more than
0.2% of (1R,3S,4R)-3-[(2'-{N,N-dimethylamino}-ethyl)]-
-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-2-one of Formula



30 or of a pharmaceutically acceptable acid addition salt
thereof.

The term "pharmaceutically acceptable acid addition
35 salts" used in the present patent specification means salts
formed with inorganic acids, e.g. hydrochloric acid,
hydrobromic acid, sulfuric acid or phosphoric acid, or

- 12 -

organic acids, e.g. acetic acid, tartaric acid, succinic acid, malic acid, lactic acid, citric acid, maleic acid or fumaric acid. The salt formed with fumaric acid possesses particularly useful properties.

5

The (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane of Formula I has three asymmetrical centres, namely in positions 1, 2, and 4.

10

According to a preferred embodiment of the invention there is provided for (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane fumarate (1 : 1), characterised by that it contains not more than 0.2% of (1R,3S,4R)-3-[(2'-{N,N-dimethylamino}-ethyl)]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-2-one fumarate (1 : 1).

15

Particularly preferably the above (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane of Formula I and pharmaceutically acceptable acid addition salts thereof according to the invention are characterised by that they contain not more than 0.1%, particularly not more than 0.05%, of the (1R,3S,4R)-3-[(2'-{N,N-dimethylamino}-ethyl)]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-2-one of Formula V or of a pharmaceutically acceptable acid addition salt thereof.

20

25


Furthermore particularly preferably the above (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane fumarate (1 : 1) is characterised by that it contains not more than 0.1%, particularly not more than 0.05%, of the (1R,3S,4R)-3-[(2'-{N,N-dimethylamino}-ethyl)]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-2-one-fumarate (1 : 1).

30

35


- 13 -

Another subject matter of the invention is a process for preparing the (1R,2S,4R)-(-)-2-[(2'-
-{N,N-dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-tri-
- [methyl]-bicyclo[2.2.1]heptane of Formula I and
5 pharmaceutically acceptable acid addition salts thereof
according to the invention by converting (+)-1,7,7-tri-
- [methyl]-bicyclo[2.2.1]heptane-2-one {(+)-camphor} of
Formula



II

into (1R,2S,4R)-(-)-2-[phenyl]-1,7,7-tri-[methyl]-
-bicyclo[2.2.1]heptane-2-ol of Formula



III

by reacting the former with a metallo-organic compound, if
necessary carrying out a decomposition, conveniently
hydrolysis, of the reaction product, and reacting the

- 14 -

(1R,2S,4R)-(-)-2-[phenyl]-1,7,7-tri-[methyl]-
-bicyclo[2.2.1]heptane-2-ol of Formula III thus obtained
with a (2-{halogeno}-ethyl)-dimethylamine in the presence
of a basic salt forming agent in an organic solvent and, if
5 desired, converting the base (1R,2S,4R)-(-)-2-[(2'-{N,N-
dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-tri-[methyl]-
-bicyclo[2.2.1]heptane of Formula I thus obtained into a
salt, characterised by carrying out the reaction of
(1R,2S,4R)-(-)-2-[phenyl]-1,7,7-tri-[methyl]-
10 -bicyclo[2.2.1]heptane-2-ol of Formula III and (2-
-{halogeno}-ethyl)-dimethylamine in a medium containing
dioxane as a solvent. The invention is not limited to the
use of dioxane as the sole solvent but also comprises the
use of a solvent containing at least 50% by weight,
15 preferably 75% by weight, of dioxane.

The essential feature of the process of the present
invention is that the alkylation is carried out in a
solvent which does not favour the alkylation reaction in
20 position 3 of (+)-camphor of Formula II in the presence of
a basic salt forming agent. It has been found that dioxane
can be used advantageously for this purpose.

Conveniently a phenyl magnesium halide is used
25 as a metallo-organic compound in a Grignard type reaction.
Further examples are phenylalkali compounds, such as
phenyllithium.

Preferably phenyl magnesium bromide is used. Phenyl
30 magnesium chloride can also be used.

35 Suitably the process according to the invention may be
carried out as follows:

- 15 -

In the first step of the process of the present invention (+)-camphor of Formula II is subjected to Grignard reaction with, for example, phenyl magnesium bromide. The reaction is carried out in a manner known per se. As reaction medium preferably tetrahydrofurane may be used. Phenyl magnesium bromide may be used in an amount of 1 to 3 moles, preferably about 1.5 mole, related to 1 mole of (+)-camphor of Formula II. One may proceed preferably by preparing first the Grignard reagent from magnesium and bromo benzene in the solvent used and thereafter adding the solution of the (+)-camphor of Formula II in an organic solvent at the boiling point of the reaction mixture. It is preferred to use the same solvent for the preparation of the Grignard reagent and the dissolving of the (+)-camphor of Formula II. As a solvent advantageously tetrahydrofurane may be used. Advantageously the reaction is carried out at the boiling point of the reaction mixture.

The reaction mixture is then cooled and the adduct obtained is hydrolysed. Hydrolysis may be carried out in a known manner, preferably in acidic medium. It is preferred to use hydrochloric acid for this purpose.

The (1R,2S,4R)-(-)-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-2-ol of Formula III obtained after decomposition of the Grignard complex can be subjected to alkylation without purifying the reaction mixture containing the same. The reaction can be carried out in the presence of unreacted (+)-camphor of Formula II. However, this leads to the formation only of a minor amount of alkylated by-products because according to the process of the present invention the formation of (1R,3S,4R)-3-[(2'-{N,N-dimethylamino}-ethyl)]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-2-one of Formula V is suppressed.

As already mentioned above, alkylation is carried out in a solvent which does not favour the alkylation reaction

- 16 -

in position 3 of (+)-camphor of Formula II, i.e. in which (+)-camphor of Formula II is alkylated in position 3 at most only to a very small extent. Dioxane is used as organic solvent because in a medium containing dioxane the alkylation of (+)-camphor of Formula II takes place at most only to a very small extent and consequently the amount of the undesired (1R,3S,4R)-3-[(2'-{N,N-dimethylamino}-ethyl)]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-2-one of Formula V in the end product (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane of Formula I is not more than 0.2% by weight.

The asymmetrical centres of (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane of Formula I in positions 1 and 4 are derived from the (+)-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-2-one {(+)-camphor} of Formula II.

The alkylation is carried out in the presence of a basic salt forming agent. The term "basic salt forming agent" means basic compounds which convert the hydroxy group into a salt. For this purpose advantageously alkali metal amides, e.g. sodium amide, or alkali metal hydrides, e.g. sodium hydride, may be used. It is preferred to use sodium amide.

Preferably as a (2-{halogeno}-ethyl)-dimethylamine (2-{chloro}-ethyl)-dimethylamine is used.

Suitably the basic salt forming agent is used in an amount of 1 to 3 moles, preferably 1.5 to 2 moles, related to 1 mole of the (1R,2S,4R)-(-)-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-2-ol of Formula III. The amount of the alkylating agent is advantageously 1.0 to 2.5 moles, preferably 1 to 1.1 mole, related to the basic salt forming agent. Preferably the alkylation reaction of

- 17 -

(1R,2S,4R)-(-)-2-[phenyl]-1,7,7-tri-[methyl]-
-bicyclo[2.2.1]heptane-2-ol of Formula III with the (2-
-{halogeno}-ethyl)-dimethylamine is carried out under
heating, particularly at the boiling point of the reaction
5 mixture. Suitably the reaction takes place within about 3
to 5 hours. An advantageous reaction time is about 4 hours.

The (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-
-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane of
10 Formula I may be converted into a pharmaceutically
acceptable salt, preferably the fumarate, optionally
without isolation of the former. One may preferably proceed
as follows: from the reaction mixture obtained after
alkylation the inorganic salts are removed by filtration at
15 0 to 30°C, preferably at 20°C, whereupon the corresponding
pharmaceutically acceptable acid, preferably fumaric acid,
is added to the filtrate in an approximately equimolar
amount (1.0 to 1.5 mole). The crystalline product
precipitated from the medium, such as the dioxane medium,
20 is filtered off.

Since the (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-
-ethoxy)]-2-[phenyl]-1,7,7-tri-[methyl]-
-bicyclo[2.2.1]heptane of Formula I or pharmaceutically
25 acceptable acid addition salts thereof, particularly the
(1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-
-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane fumarate
(1 : 1) obtained by the process according to the invention
contain[s] in accordance with the requirements of
30 Pharmacopoeia not more than 0.2% of (1R,3S,4R)-3-[(2'-{N,N-
-dimethylamino}-ethyl)]-1,7,7-tri-[methyl]-
-bicyclo[2.2.1]heptane-2-one of Formula V or of the
pharmaceutically acceptable acid addition salt thereof,
respectively, by the process according to the invention
35 recrystallization from dimethyl formamide anyhow leading
only to insufficient purification used by known methods has
been eliminated, and thus there is no need to remove traces

- 18 -

of dimethyl formamide from the pharmaceutically active principle by methods unsuitable for the given purpose, too. Also this latter is a significant progress in view of the impossibility to remove dimethyl formamide to the necessary extent because of its high boiling temperature at which
5 (1R,2S,4R) - (-) - 2 - [(2' - {N,N-dimethylamino} - ethoxy)] - 2 - [phenyl] - 1,7,7-tri - [methyl] - bicyclo[2.2.1]heptane of Formula I would be decomposed.

10 The advantage of the process of the present invention is that in addition to providing a highly pure product meeting the strict requirements of Pharmacopoeia it can be carried out with excellent yields. Thus the yield of about 46% shown in the Examples is considerably higher than the
15 yields disclosed in the Prior Art which do not surpass 25% even if (+) - camphor is re-circulated several times.

A further subject matter of the invention are medicaments characterised by that they contain as [an]
20 principle(s) 1 or more compound(s) according to the invention within the above definition, advantageously together with 1 or more in pharmaceutical preparations usual excipient(s).

25 Preferably the medicaments according to the invention contain (1R,2S,4R) - (-) - 2 - [(2' - {N,N-dimethylamino} - ethoxy)] - 2 - [phenyl] - 1,7,7-tri - [methyl] - bicyclo[2.2.1]heptane fumarate (1 : 1) according to the invention within the above definition as an active principle.

30 Suitably the medicaments according to the invention are in the form of pharmaceutical preparations. They may be prepared by known methods of the pharmaceutical technique. The preparations are preferably suitable for oral
35 administration, e.g. tablets, coated tablets, capsules, solutions, emulsions or suspensions, or for parenteral administration, e.g. intravenous, percutaneous or

- 19 -

intramuscular injectable solutions. The compositions may contain usual carriers, e.g. starch, lactose and/or calcium carbonate, and/or water, polyalkylene glycols, sodium chloride solution and/or dextrose solution. The pharmaceutical preparations may also contain usual pharmaceutical auxiliary agents, e.g. emulsifying, stabilising, suspending and/or disintegrating agents, salts to modify the osmotic pressure, buffers and/or antioxidants.

A still further subject matter of the invention is the use of the compounds according to the invention within the above definition for preparing anxiolytic medicaments.

Preferably (1R,2S,4R) - (-) - 2 - [(2' - {N,N-dimethylamino} - ethoxy)] - 2 - [phenyl] - 1,7,7-tri - [methyl] - bicyclo[2.2.1]heptane fumarate (1 : 1) is used as a compound according to the invention within the above definition.

The invention is further illustrated by the following Examples. The melting points given in the Examples are uncorrected values.

Example 1

(1R,2S,4R) - (-) - 2 - [(2' - {N,N-dimethylamino} - ethoxy)] - 2 - [phenyl] - 1,7,7-tri - [methyl] - bicyclo[2.2.1]heptane fumarate (1 : 1) [Formula I]

Grignard reaction

To a suspension of 48.6 g (1.5 g atom) magnesium spans and 600 ml of anhydrous tetrahydrofurane a 20 ml portion of a mixture of 236 g (1.5 moles) of bromo benzene and 200 ml of anhydrous tetrahydrofurane is added at the boiling

- 20 -

point. Once the Grignard reaction has started, the residual part of the bromo benzene mixture is added to the suspension dropwise within an hour. The reaction mixture is heated to boiling until the magnesium is completely dissolved. To the Grignard compound a solution of 152.2 g (1.0 mole) of (+)-camphor of Formula II and 300 ml of anhydrous tetrahydrofuran is added under constant heating to boiling within about half an hour and the reaction mixture is heated to boiling for a further period of 5 hours.

Hydrolysis

The reaction mixture is cooled to 25°C and poured onto a mixture of 500 ml of heptane, 400 g of ice, 30 g of sodium chloride and 150 ml of concentrated hydrochloric acid under stirring at 0°C. The organic phase is separated and made alkaline to pH 10 by adding a 25% by weight/volume aqueous ammonium hydroxide solution. After repeated separation the solution is dried and evaporated in vacuo. Thus 220 g of a colourless oil are obtained.

Analysis on the basis of GC

- The test is carried out on a Perkin Elmer Autosystem gas chromatograph.
- Length 10 m (0.25 mm).
- A 14% cyanopropyl 14% methyl polysiloxane fixed phase (CPSil-19CB, Chrompack [Handelsprodukt]) capillary column is used.
- Injection is performed at 200°C.
- Heating speed 10°C/minute.
- Carrier gas: helium.
- Detector: FID, injection temperature 200°C, final temperature 250°C, gas pressure 40 kPa.

- 21 -

(1R,2S,4R)-(-)-2-[phenyl]-1,7,7-tri-[methyl]-
-bicyclo[2.2.1]heptane-2-ol content: 66.5%

(+)-camphor content: 25%.

5

Etherification

To a suspension of 45.5 g (1.05 mole) sodium amide
(content: 90% by weight/weight) and 500 ml of anhydrous
10 dioxane a mixture of 220 g of the colourless oil obtained
by the hydrolysis containing (1R,2S,4R)-(-)-2-[phenyl]-
-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-2-ol, and 100 ml
of anhydrous dioxane is added at the boiling point within
half an hour. The mixture is heated to boiling for 2 hours,
15 whereupon 113.0 g (1.05 mole) of (2-{chloro}-ethyl)-
-dimethylamine are added and the reaction mixture is heated
to boiling for a further period of 4 hours.

Formation of the fumarate salt

20

The suspension is cooled to 20°C, filtered, to the
clear filtrate 121.9 g (1.05 mole) of fumaric acid are
added under vigorous stirring. The reaction mixture is
heated to boiling for 10 minutes, cooled to 15°C, stirred
25 for a further period of an hour and filtered. The filter
cake is washed with dioxane, water and ethanol and dried at
80°C until free of solvent. Thus 190.5 g (0.456 mole) of
white crystals are obtained, yield 45.6% [based on
(+)-camphor]. The melting point of the white crystals
30 amounts to 214 to 216°C.

Analysis for the Formula $C_{20}H_{31}NO.C_4H_4O_4$ (417.55)

35	calculated:	C% = 69.03%;	H% = 8.45%;	N% = 3.35%;
	found:	C% = 69.06%;	H% = 8.42%;	N% = 3.39%.

$[\alpha]_D^{20} = -92.5^\circ$ (c = 0.4, dimethyl sulfoxide, 435 nm).

- 22 -

The product contains less than 0.05% of (1R,3S,4R)-3-
-[(2'-{N,N-dimethylamino}-ethyl)]-1,7,7-tri-[methyl]-
-bicyclo[2.2.1]heptane-2-one fumarate (1 : 1)
contamination.

5

Example 2 (Comparative Example)

10

(1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-
-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane
fumarate (1 : 1) [Formula I]

Reproduction of the process disclosed in Hungarian
patent No. 212,574

15

Grignard reaction

20

To a Grignard compound prepared from 5.52 g (0.23 g
atom) of magnesium spans and 36.1 g (0.23 mole) of bromo
benzene in 200 ml of anhydrous diethyl ether a solution of
30.4 g (0.20 mole) of (+)-camphor and 50 ml of anhydrous
diethyl ether is added. The reaction mixture is heated to
boiling for 5 hours. The Grignard complex is decomposed by
adding an icecold aqueous solution of 20 g of ammonium
chloride, the mixture is washed three times with 30 ml of
water each, separated, dried over anhydrous magnesium
sulfate and the solvent is removed by evaporation. Thus
40.5 g of a colourless oil are obtained which contains
according to GC

30

57.5% of (+)-camphor of Formula II
5.8% of 1,7,7-tri-[methyl]-
-bicyclo[2.2.1]heptane-2-ol [borneol];
34.5% of (1R,2S,4R)-(-)-2-[phenyl]-1,7,7-tri-
- [methyl]-bicyclo[2.2.1]heptane-2-ol
of Formula III; and

35

- 23 -

2.2% of further contaminations in smaller amounts.

Etherification

5

To a suspension of 3.4 g (67 millimoles) of sodium hydride (47.5% by weight/weight dispersion) and 50 ml of anhydrous toluene a solution of 40.0 g of the oil obtained in the Grignard reaction containing (1R,2S,4R)-(-)-2-[phenyl]-1,7,7-tri-
10 -[methyl]-bicyclo[2.2.1]heptane-2-ol of Formula III and 30 ml of anhydrous toluene are added. The reaction mixture is heated to boiling for an hour, whereupon a solution of 6.85 g (67 millimoles) of (2-{chloro}-ethyl)-
15 -dimethylamine and 10 ml of toluene is added at the boiling point. The reaction mixture is heated to boiling for a further period of 4 hours.

Separation

20

The reaction mixture is washed three times with 25 ml of water each. The product is extracted with three equal portions of a solution of 18 g (0.12 mole) of tartaric acid and 40 ml of water. The phases are separated, the aqueous
25 layers are combined, made alkaline to pH 10 with a concentrated ammonium hydroxide solution, extracted three times with 20 ml of dichloro ethane each, dried over magnesium sulfate and the solvent is removed in vacuo. Thus 14.5 g of a colourless oil are obtained which contains
30 according to GC analysis

74.2% of (1R,2S,4R)-(-)-2-[(2'-{N,N-
-dimethylamino}-ethoxy)]-2-[phenyl]-
-1,7,7-tri-[methyl]-
35 -bicyclo[2.2.1]heptane of Formula I;

- 24 -

16.5% of (1R,4R)-2-[(2'-{N,N-dimethylamino}-
-ethoxy)]-1,7,7-tri-[methyl]-
-bicyclo[2.2.1]heptane of Formula IV;

5 6.5% of (1R,3S,4R)-3-[(2'-{N,N-dimethylamino}-
-ethyl)]-1,7,7-tri-[methyl]-
-bicyclo[2.2.1]heptane-2-one of Formula V;
and
some % of further unidentified contaminations each
10 in an amount below 1%.

Formation of the fumarate salt

To a solution of 14.0 g of the base (1R,2S,4R)-(-)-2-
15 -[(2'-{N,N-dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-tri-
-[methyl]-bicyclo[2.2.1]heptane of Formula I set free from
the tartrate salt and 150 ml of ethanol 5.07 g (43.6
millimol) of fumaric acid are added at 70°C. The product is
filtered at 0°C and recrystallized from 50 ml of dimethyl
20 formamide.

Thus 13.5 g of the desired product (1R,2S,4R)-(-)-2-
-[(2'-{N,N-dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-tri-
-[methyl]-bicyclo[2.2.1]heptane fumarate (1 : 1) of Formula
25 I are obtained in the form of white crystals. Yield 16.2%
[based on (+)-camphor]. According to GC analysis in the
product 0.5% of (1R,3S,4R)-3-[(2'-{N,N-dimethylamino}-
-ethyl)]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-2-one of
Formula V can be detected. Mp.: 214 to 216°C.

30

Analysis for the Formula $C_{20}H_{31}NO.C_4H_4O_4$ (417.55)


calculated: C% = 69.03%; H% = 8.45%; N% = 3.35%;

35


found: C% = 69.16%; H% = 8.52%; N% = 3.32%.

Claims

- 1.) (1R,2S,4R) - (-) - 2 - [(2' - {N,N-dimethylamino} - ethoxy)] - 2 -
- [phenyl] - 1,7,7-tri - [methyl] - bicyclo[2.2.1]heptane of
Formula



and pharmaceutically acceptable acid addition salts
thereof, characterised by that they contain not more
than 0.2% of (1R,3S,4R) - 3 - [(2' - {N,N-dimethylamino} -
ethyl)] - 1,7,7-tri - [methyl] - bicyclo[2.2.1]heptane-2-
-one of Formula




or of a pharmaceutically acceptable acid addition salt
thereof.

- 26 -

- 2.) (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-
- [phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane
fumarate (1 : 1) according to claim 1, characterised
by that it contains not more than 0.2% of (1R,3S,4R)-
5 -3-[(2'-{N,N-dimethylamino}-ethyl)]-1,7,7-tri-
- [methyl]-bicyclo[2.2.1]heptane-2-one fumarate
(1 : 1).
- 3.) (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-
10 - [phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane of
Formula I according to claim 1 or 2, and
pharmaceutically acceptable acid addition salts
thereof, characterised by that they contain not more
than 0.05% of the (1R,3S,4R)-3-[(2'-{N,N-dimethyl-
15 amino}-ethyl)]-1,7,7-tri-[methyl]-
-bicyclo[2.2.1]heptane-2-one of Formula V or of a
pharmaceutically acceptable acid addition salt
thereof.
- 20 4.) (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-
- [phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane
fumarate (1 : 1) according to any of claims 1 to 3,
characterised by that it contains not more than 0.05%
of the (1R,3S,4R)-3-[(2'-{N,N-dimethylamino}-
25 -ethyl)]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-2-
-one fumarate (1 : 1).
- 5.) Process for preparing the (1R,2S,4R)-(-)-2-
- [(2'-{N,N-dimethylamino}-ethoxy)]-2-[phenyl]-1,7,7-
30 -tri-[methyl]-bicyclo[2.2.1]heptane of Formula I and
pharmaceutically acceptable acid addition salts
thereof according to any of claims 1 to 4 by
converting (+)-1,7,7-tri-[methyl]-
-bicyclo[2.2.1]heptane-2-one {(+)-camphor} of Formula
35

- 27 -

5




II

10

into (1R,2S,4R)-(-)-2-[phenyl]-1,7,7-tri-[methyl]-
-bicyclo[2.2.1]heptane-2-ol of Formula

15



III

20

25

by reacting the former with a metallo-organic
compound, if necessary carrying out a decomposition of
the reaction product, and reacting the (1R,2S,4R)-(-)-
-2-[phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane-
-2-ol of Formula III thus obtained with a (2-
-{halogeno}-ethyl)-dimethylamine in the presence
of a basic salt forming agent in an organic solvent
and, if desired, converting the

30

35

base (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-
-ethoxy)]-2-[phenyl]-1,7,7-tri-[methyl]-
-bicyclo[2.2.1]heptane of Formula I thus obtained into
a salt, characterised by carrying out the reaction of
5 (1R,2S,4R)-(-)-2-[phenyl]-1,7,7-tri-[methyl]-
-bicyclo[2.2.1]heptane-2-ol of Formula III and (2-
-{halogeno}-ethyl)-dimethylamine in a medium
containing dioxane as solvent.

- 10 6.) Process according to claim 5, characterised by that a
phenyl magnesium halide is used as a metallo-
-organic compound in a Grignard type reaction.
- 15 7.) Process according to claim 5 or 6, characterised by
that phenyl magnesium bromide is used.
- 20 8.) Process according to any of claims 5 to 7,
characterised by that phenyl magnesium chloride
is used.
- 25 9.) Process according to any of claims 5 to 8,
characterised by that as a (2-{halogeno}-ethyl)-
-dimethylamine (2-{chloro}-ethyl)-dimethylamine is
used.
- 30 10.) Process according to any of claims 5 to 9,
characterised by that as a basic salt forming agent
sodium amide is used.
- 35 11.) Process according to any of claims 5 to 10,
characterised by carrying out the alkylation reaction
of (1R,2S,4R)-(-)-2-[phenyl]-1,7,7-tri-[methyl]-
-bicyclo[2.2.1]heptane-2-ol of Formula III with the
(2-{halogeno}-ethyl)-dimethylamine under heating,
particularly at the boiling point of the reaction
mixture.

- 29 -

12.) Process according to any of claims 5 to 11,
characterised by that the conversion of the
(1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-
- [phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane of
5 Formula I into the fumarate (1 : 1) is carried out
without isolation of the former.

13.) Medicaments, characterised by that they contain as
[an] active principle(s) 1 or more compound(s)
10 according to any of claims 1 to 4 within the
definitions there, advantageously together with 1 or
more in pharmaceutical preparations usual
excipient(s).

14.) Medicaments according to claim 13, characterised by
that they contain (1R,2S,4R)-(-)-2-[(2'-{N,N-dimethyl-
amino}-ethoxy)]-2-[phenyl]-1,7,7-tri-[methyl]-
-bicyclo[2.2.1]heptane fumarate (1 : 1) according to
claim 2 or 4 as an active principle.

15.) The use of the compounds according to any of claims 1
to 4 within the definitions there for preparing
anxiolytic medicaments.

16.) The use according to claim 15, characterised by that
(1R,2S,4R)-(-)-2-[(2'-{N,N-dimethylamino}-ethoxy)]-2-
- [phenyl]-1,7,7-tri-[methyl]-bicyclo[2.2.1]heptane
fumarate (1 : 1) of claim 2 or 4 is used as a compound
according to claims 1 to 4.